

JOURNAL OF ALGEBRA 8, 156-164 (1968)

The Structure of Certain Artinian Rings with Zero Singular Ideal

R. R. COLBY AND EDGAR A. RUTTER, JR.

The University of Kansas, Lawrence, Kansas 66044

Communicated by A. W. Goldie

Received March 4, 1967

INTRODUCTION

Let R be a ring with identity and M and N (left) R -modules with $M \subset N$. Then M is said to be *essential* in N if M intersects every nonzero submodule of N nontrivially. A left ideal I of R is called an essential left ideal if I is essential in R . We denote the *singular submodule* of N by $Z(N)$; $Z(N) = \{n \in N \mid In = (0) \text{ for some essential left ideal } I \text{ of } R\}$. We shall assume throughout this paper that $Z(R) = 0$. In Section 1 we outline the construction of Johnson's ring of quotients [3] and some of its properties and develop certain other preliminary results. In Section 2 we determine the structure of Artinian rings with zero singular ideal having the property that each principal indecomposable left ideal contains a unique minimal left ideal. In particular we show that if R is such a ring then R is a certain distinguished subring of a complete blocked triangular matrix ring over a division subring of R . We also consider the case where R is a left generalized uniserial ring and generalize results of A. W. Goldie [2] and I. Murase [5]. Finally we apply the methods to obtain the structure of finite-dimensional algebras with zero singular ideal.

1. PRELIMINARIES

If R is a ring with zero singular ideal, then Johnson [3] has constructed a ring extension Q of R which we shall call Johnson's ring of quotients of R . Essentially, the definition of Q in [3] was as follows: $Q = \bigcup_A \text{Hom}_R(A, R)$, where A is an essential left ideal of R . If $x, y \in Q$, then we take $x = y$ if and only if $xa = ya$ for every a in some essential left ideal

$$A \subset (\text{domain } x) \cap (\text{domain } y).$$

This extension Q of R is a ring of quotients of R in the sense that ${}_R Q$ is an

essential extension of ${}_R R$ and since $Z({}_R R) = 0$ so does $Z({}_R Q)$. The following property of Q is established by Lambek [4].

LEMMA 1.1. *If I and J are left ideals of R with I essential in R , then $\text{Hom}_R(I, J)$ is isomorphic to*

$$J : I = \{q \in Q \mid Iq \subset J\}$$

under the correspondence $(i \rightarrow iq) \leftrightarrow q, q \in Q, i \in I$.

LEMMA 1.2. *Suppose that M and N are submodules of ${}_R Q$ and $f \in \text{Hom}_R(M, N)$. Then there exists a unique $\bar{f} \in \text{Hom}_Q(QM, QN)$ such that $\bar{f}|_M = f$. If f is one-to-one or onto, so is \bar{f} .*

Proof. For $x = \sum q_i m_i \in QM$, define $\bar{f}(x) = \sum q_i f(m_i)$. The assertions of the lemma are easily verified.

We denote the left and right annihilators of a subset $S \subset R$ by S_l and S_r , respectively.

LEMMA 1.3. *If I is a left ideal of R , then $I_r = \{x \in R \mid x_l \cap I \text{ is essential in } I\}$.*

Proof. $x \in I_r \Rightarrow x_l \supset I \Rightarrow x_l \cap I$ is essential in I . Conversely, if $x_l \cap I$ is essential in I , then $(x_l \cap I : y)yx = 0$ for all $y \in I$. But $(x_l \cap I : y)$ is essential in R . For, if $(x_l \cap I : y) \cap I' = 0$, then $I'y \cap (x_l \cap I) = 0$ so $I'y = 0$. Thus $I' \subset (x_l \cap I : y) \cap I' = 0$ and $I' = 0$. Now since $(x_l \cap I : y)yx = 0$, we get $yx = 0$. Hence $x \in I_r$.

We say that M is a *uniform* R -module if every nonzero submodule of M is an essential submodule of M .

LEMMA 1.4. *If M is a uniform R -module, I is a left ideal of R and $0 \neq \alpha \in \text{Hom}_R(M, I)$, then α is one-to-one.*

Proof. If $\ker \alpha \neq 0$ then for any $m \in M$, $(\ker \alpha : m)$ is an essential left ideal of R . Furthermore,

$$(\ker \alpha : m) \alpha(m) = \alpha(\ker \alpha m) = 0$$

and hence $\alpha(m) = 0$ since $Z(R) = 0$. Thus $\alpha = 0$, a contradiction.

LEMMA 1.5 (Dual Bases Lemma, [1], Chapter VII, Prop. 3.1). *An R -module M is projective if and only if there exists a family $\{a_\alpha\} \subset M$ and a family $\{\lambda_\alpha\} \subset \text{Hom}_R(M, R)$ such that, for each $m \in M$, $m = \sum_\alpha \lambda_\alpha(m) a_\alpha$ where $\lambda_\alpha(m) = 0$ for all but a finite number of indices α .*

For the rest of this section we shall assume that R is left Artinian. We denote the (left) socle of R by E . Then E is the unique minimal essential left ideal of R .

LEMMA 1.6. *The socle E of R is a projective left R -module.*

Proof. Let M be a minimal left ideal of R . Since $EM \neq 0$, there exists a minimal left ideal $I \subset E$ such that $IM \neq 0$ and hence $IM = M$. Since I is not nilpotent, $I = Re$ with $e^2 = e$ and $M \cong I$ is projective. Thus E is clearly projective.

PROPOSITION 1.7. $Q = QE$.

Proof. Since E is projective, Lemma 1.5 implies that there exists $\{a_\alpha\} \subset E$ and $\{\lambda_\alpha\} \subset \text{Hom}_R(E, R)$ such that

$$y = \sum_{\alpha} \lambda_{\alpha}(y) a_{\alpha}$$

for all $y \in E$. By Lemma 1.1 there exist $q_{\alpha} \in Q$ such that $\lambda_{\alpha}(y) = yq_{\alpha}$ for all $y \in E$. Hence $y = \sum_{\alpha} yq_{\alpha}a_{\alpha}$ and so $E(1 - \sum_{\alpha} q_{\alpha}a_{\alpha}) = 0$. Thus, since $Z(R) = 0$, $1 = \sum_{\alpha} q_{\alpha}a_{\alpha} \in QE$.

2. THE STRUCTURE THEOREM

We shall assume throughout the rest of this paper that R is a left Artinian ring with zero singular ideal having the property that every principal indecomposable left ideal contains a unique minimal left ideal.

LEMMA 2.1. *If R has the above properties, then R can be written as a direct sum of indecomposable (two-sided) ideals with the above properties as rings.*

Proof. Let e_i be primitive idempotents and let M_i be the unique minimal left ideal of Re_i for $i = 1, 2$. If $M_1 \not\cong M_2$, then we must have $M_1Re_2 = 0$ and hence it follows from Lemma 1.3 that $Re_1Re_2 = 0$. Similarly, $Re_2Re_1 = 0$. Thus if we write R as a direct sum of indecomposable left ideals and group these according to the isomorphism type of their unique minimal ideals, we clearly obtain a decomposition of R as a direct sum of ideals which as rings have the above properties. Since each of these rings clearly has homogeneous socle they are indecomposable.

We shall assume henceforth that R is indecomposable.

LEMMA 2.2. *The irreducible left ideals of R are R -isomorphic and R has an irreducible summand.*

Proof. This is immediate from the reasoning of the above lemma and Lemma 1.6.

Let e_1, \dots, e_m be a complete set of primitive idempotents of R . We say that $e_i \sim e_j$ if and only if $Re_i \not\cong Re_j$, for all $1 \leq i, j \leq m$. This is clearly an equivalence relation. We partially order the resulting equivalence classes by saying that $[e_i] \leq [e_j]$ if and only if Re_i is isomorphic to a submodule of Re_j . This is a partial order since R is left-Artinian. Note that this partially ordered set contains a unique minimal element, namely, the equivalence class determined by the idempotent corresponding to an irreducible summand of R .

LEMMA 2.3. *$[e_i] \leq [e_j]$ if and only if $e_i Re_j \neq 0$.*

Proof. The necessity is obvious and the sufficiency follows easily from Lemma 1.4.

LEMMA 2.4. *Let e_1, \dots, e_m be a complete set of primitive orthogonal idempotents of R . Then there exists an integer k , k integers n_1, n_2, \dots, n_k with $\sum_{i=1}^k n_i = m$ and an indexing e_j^i of the above idempotents with $i = 1, \dots, k$ and $1 \leq j \leq n_i$ such that $e_j^i \sim e_{j'}^{i'}$, for all i, j , and j' and $e_j^i Re_j^i = 0$ if $i' > i$, for all j and j' .*

Proof. Let k be the number of equivalence classes of idempotents resulting from the above equivalence relation. First number the equivalence classes from 1 to k in such a way that if $[e_j]_{i'} \leq [e_j]_i$ then $i' \geq i$. Now let n_i be the number of idempotents in the i th class. If we index the idempotents of the i th class with a superscript i and a subscript ranging from 1 to n_i , then it is immediate from the definition of \sim and Lemma 2.3 that this is the desired indexing.

LEMMA 2.5. *Re_j^i is irreducible if and only if $i = k$.*

Proof. This is immediate from Lemma 2.2 and the manner in which the idempotents were indexed above.

LEMMA 2.6. *For all i, j , $e_j^i Re_j^i$ is a division ring.*

Proof. Let N denote the radical of R . It suffices to show that $e_j^i Ne_j^i = 0$. If this were not so then Lemma 1.4 would imply that Re_j^i is isomorphic to a submodule of Ne_j^i .

LEMMA 2.7. $Q = Qe_1^1 \oplus \cdots \oplus Qe_{n_1}^1 \oplus \cdots \oplus Qe_1^k \oplus \cdots \oplus Qe_{n_k}^k$, each Qe_j^i is an irreducible left ideal in Q and $Qe_j^i \cong Qe_{j'}^{i'}$ for all i, i', j , and j' . In particular Q is a simple ring.

Proof. Let M_j^i denote the unique irreducible submodule of Re_j^i . Then by Proposition 1.7,

$$\begin{aligned} Q &= QE \\ &= QM_1^1 \oplus \cdots \oplus QM_{n_1}^1 \oplus \cdots \oplus QM_1^k \oplus \cdots \oplus QM_{n_k}^k \\ &\subset Qe_1^1 \oplus \cdots \oplus Qe_{n_1}^1 \oplus \cdots \oplus Qe_1^k \oplus \cdots \oplus Qe_{n_k}^k = Q. \end{aligned}$$

Hence $QM_j^i = Qe_j^i$ for each i, j . Since $M_j^i \cong_R M_{j'}^{i'}$, Lemma 1.2 implies that $Qe_j^i \cong_Q Qe_{j'}^{i'}$. Also, since ${}_R M_j^i$ is essential in ${}_R(Qe_j^i)$ and ${}_R M_j^i$ is irreducible, M_j^i is contained in every nonzero Q -submodule of Qe_j^i . Thus every nonzero Q -submodule of Qe_j^i contains $QM_j^i = Qe_j^i$. Thus Qe_j^i is an irreducible Q -module.

LEMMA 2.8.

$$\text{Hom}_Q(Qe_1^k, Qe_1^k) \cong e_1^k Qe_1^k = e_1^k Re_1^k \cong \text{Hom}_R(Re_1^k, Re_1^k).$$

Proof. Since Re_1^k is irreducible, it is the socle of Qe_1^k . Thus $Re_1^k qe_1^k \subset Re_1^k$, for all $q \in Q$. Thus there exists $r \in R$ such that $e_1^k r e_1^k = e_1^k q e_1^k$.

THEOREM 2.9. *There exists a positive integer k , and positive integers n_1, n_2, \dots, n_k such that R is isomorphic to a blocked triangular matrix ring such that the i, j th block is an arbitrary $n_i \times n_j$ matrix with entries in an additive subgroup D_{ij} of a fixed division subring D of R . These subgroups satisfy the following additional properties:*

- (1) $D_{il}D_{lj} \subset D_{ij}$, for all i, j, l .
- (2) D_{ii} is a division ring for each i .
- (3) $D_{ki} = D$ for all i .
- (4) D_{ij} is finite dimensional as a left vector space over D_{ii} for all i and j .

Conversely, if D is a fixed division ring, k and n_1, \dots, n_k fixed integers and D_{ij} are fixed subgroups of D with the above properties then the corresponding blocked triangular matrix ring is a left Artinian ring with zero singular ideal and such that each principal indecomposable left ideal contains a unique minimal left ideal.

Proof. Let e_1, \dots, e_m be a complete set of primitive orthogonal idempotents for R , k a positive integer, n_1, \dots, n_k positive integers, and e_j^i an indexing of these idempotents as described in Lemma 2.4. Let g_t^i be a fixed isomorphism of Re_t^i onto Re_1^i and let

$$r_t^i = g_t^i(e_t^i) \in e_t^i Re_1^i,$$

and

$$s_t^i = (g_t^i)^{-1}(e_1^i) \in e_1^i Re_t^i.$$

A computation shows that

$$r_t^i s_t^i = e_t^i \quad \text{and} \quad s_t^i r_t^i = e_1^i. \quad (*)$$

Let h^i be a fixed isomorphism of Qe_1^i onto Qe_1^k with h^k equal the identity on Qe_1^k ,

$$\alpha^i = h^i(e_1^i) \in e_1^i Qe_1^k,$$

and

$$\beta^i = (h^i)^{-1}(e_1^k) \in e_1^k Qe_1^i.$$

A computation shows that

$$\alpha^i \beta^i = e_1^i \quad \text{and} \quad \beta^i \alpha^i = e_1^k. \quad (**)$$

We define a function θ from R into the ring of $m \times m$ blocked triangular matrices with entries in $D = e_1^k Re_1^k$ by letting $\theta(a)$ be the matrix whose (i, j) th block is the $n_i \times n_j$ matrix

$$A_{ij} = (\beta^i s_u^i e_u^i a e_v^j r_v^j \alpha^j) = (\beta^i s_u^i a r_v^j \alpha^j).$$

The entries in this block belong to

$$D_{ij} = \{\beta^i x \alpha^j \mid x \in R\},$$

which is an additive subgroup of $e_1^k Qe_1^k = e_1^k Re_1^k$, isomorphic to $e_1^i Re_1^j$. Since this is a ring isomorphism when $i = j$, D_{ii} is isomorphic to $e_1^i Re_1^i$ which is a division ring.

We will now show that θ maps onto the blocked triangular matrix ring whose (i, j) th block has arbitrary entries in D_{ij} by showing that there exists an element $\bar{x} \in R$ such that $(A_{ij})_{uv} = \beta^i x \alpha^j$ for any i, j, u, v and any $x \in R$, and all other entries of $\theta(\bar{x})$ are zero. Let $\bar{x} = r_u^i x s_v^j \in e_u^i Re_v^j$. Then since the $e_{j'}^i$ are orthogonal idempotents, all the entries of $\theta(\bar{x})$ are zero except the (u, v) th entry of the (i, j) th block, while

$$\begin{aligned} (A_{ij})_{uv} &= \beta^i s_u^i r_u^i x s_v^j r_v^j \alpha^j \\ &= \beta^i e_1^i x e_1^j \alpha^j = \beta^i x \alpha^j. \end{aligned}$$

Clearly from the choice of h^k , $\alpha^k = \beta^k = e_1^k$ and so $D_{kk} = e_1^k R e_1^k = D$. Also $D_{kj} \subset D_{kk}$. But for any j , $e_1^k x \beta^j \in e_1^k R e_1^k Q e_1^j \subset e_1^k R e_1^k Q \subset EQ$ since $R e_1^k$ is an irreducible R -module (Lemma 2.5). But $EQ \subset R$ since ${}_R Q$ is an essential extension of ${}_R R$ and E is the unique minimal essential left ideal of R . Thus $e_1^k x \beta^j \in R$ and so $e_1^k x \beta^j \alpha^j = e_1^k x e_1^k \in D_{kj}$. Thus $D_{kj} \supset D_{kk}$.

One computes that θ is a ring homomorphism and that $D_{ii} D_{ij} \subset D_{ij}$ for all i and j using conditions (*) and (**). For any $0 \neq a \in R$, $e_j^i a e_{j'}^{i'} \neq 0$ for some i, i', j, j' , and hence it follows from (*) and (**) that $\theta(a) \neq 0$. Thus θ is one-to-one.

We have, therefore, established an isomorphism between R and a ring of matrices of the type described in the theorem and have verified the necessity of (1)-(3). Condition (4) is most easily verified by noting that the matrix ring and hence also R would not be left-Artinian if (4) did not hold.

The converse of the theorem follows readily from an examination of matrix rings of the type described above.

We say that R is *left generalized uniserial* if each principal indecomposable left ideal of R has a unique composition series.

COROLLARY 2.10. *R is an (indecomposable) left generalized uniserial ring (with zero left singular ideal) if and only if R has a matrix representation of the type described in the preceding theorem which has the following additional properties:*

- (1) D_{ij} has dimension at most one as a (left) vector space over D_{ii} for all i and j .
- (2) If $D_{ij} \neq 0$ and $D_{i'j} \neq 0$ with $i \leq i'$, then $D_{i'i} \neq 0$.

Proof. A left generalized uniserial ring satisfies the hypothesis of Theorem 2.9 and hence has a matrix representation of the type described there. That (1) and (2) are necessary and sufficient for such a matrix ring to be left generalized uniserial follows from a straightforward examination of these matrix rings.

We say that R is *meet irreducible* if and only if any two nonzero ideals of R have nonzero intersection.

COROLLARY 2.11. *R is a meet irreducible left generalized uniserial ring (with zero left singular ideal) if and only if R has a matrix representation of the type described in Theorem 2.9 with the additional property that D_{ij} is one-dimensional as a left vector space over D_{ii} for all i and j with $j \leq i$.*

LEMMA 2.12. *Let A be a finite-dimensional algebra over a field K . Then A has zero left singular ideal if and only if A is isomorphic to a subalgebra of a finite-dimensional K -algebra \hat{A} with the following properties:*

(1) *Each principal indecomposable left ideal of \hat{A} contains a unique minimal left ideal.*

(2) *\hat{A} has zero left singular ideal.*

(3) *The socles of ${}_A A$ and ${}_A \hat{A}$ coincide.*

Proof. Let Q be Johnson's (left) ring of quotients for A . Johnson [3] has shown that Q is a semisimple ring with minimum condition. This can also be seen by modifying slightly the proof of Lemma 2.7. We note that Q is a finite-dimensional algebra over K since ${}_A Q$ is the injective envelope of ${}_A A$.

Let e_1, \dots, e_m be a complete set of primitive orthogonal idempotents for A . We can write each e_i as

$$e_i = e_i^1 + \dots + e_i^{k_i},$$

so that the

$$\{e_i^j \mid j = 1, \dots, k_i; i = 1, \dots, m\}$$

is a complete set of primitive orthogonal idempotents for Q . Note that k_i is the number of summands in any irreducible decomposition of the socle of Ae_i . For, let $M_1^i \oplus \dots \oplus M_{u_i}^i$ be the socle of Ae_i . Then, by Proposition 1.6,

$$\begin{aligned} Q &= QE = QM_1^1 \oplus \dots \oplus QM_{u_1}^1 \oplus \dots \oplus QM_1^m \oplus \dots \oplus QM_{u_m}^m \\ &= Qe_1 \oplus \dots \oplus Qe_m. \end{aligned}$$

Hence $Qe_i = QM_1^i \oplus \dots \oplus QM_{u_i}^i$ and so $u_i = k_i$ since the QM_j^i are irreducible Q -modules. Thus the socle of A contains $t = \sum_{i=1}^m k_i$ components. Note also that Ae_i contains a unique minimal submodule if and only if $k_i = 1$.

Let \hat{A} be the smallest subalgebra of Q which contains A and $\{e_i^j \mid j = 1, \dots, k_i; i = 1, \dots, m\}$. Since Q is a finite-dimensional K -algebra so is \hat{A} . Let I be an irreducible left ideal of A . Then ${}_A I$ is projective by Lemma 1.6 and so $e_i I \neq 0$ implies that Ae_i is irreducible. Suppose $e_i^j I \neq 0$. Then $0 \neq e_i^j I = e_i^j e_i I$ and so Ae_i is an irreducible projective, $k_i = 1$, and $e_i^j = e_i$. Thus $e_i^j I \subset I$. Now any element of \hat{A} is a sum of products of terms of the form ae_i^j with $a \in A$ and so $\hat{A}I \subset I$. Thus irreducible submodules of ${}_A A$ are irreducible submodules of ${}_A \hat{A}$. Furthermore, for all i, j , $\hat{A}e_i^j$ is an A -submodule of Qe_i^j and hence contains a unique minimal left ideal of A , and hence also a unique minimal left ideal of \hat{A} . Since there are t irreducible components in the socle of ${}_A A$, the socles of ${}_A A$ and ${}_A \hat{A}$ must coincide. Thus since ${}_A \hat{A}$ is an essential extension of ${}_A A$, ${}_A \hat{A}$ has zero left singular ideal. The converse is immediate.

THEOREM 2.13. *Let A be a finite-dimensional algebra over a field K . Then*

A has zero left singular ideal if and only if A is (isomorphic to) a subalgebra of an algebra T with the following properties:

(1) $T = T_1 \oplus \cdots \oplus T_n$, where each T_i is a two sided ideal of T and T_i is a complete blocked triangular matrix ring over a finite dimensional division algebra D_i over K .

(2) The socles of ${}_A A$ and ${}_T T$ coincide.

Proof. This theorem is immediate from Lemma 2.12 and Theorem 2.9.

Remark. We have used the hypothesis that A is a finite-dimensional algebra in the proof of Lemma 2.12 only to show that \hat{A} is a ring with left minimum condition. Thus Theorem 2.13 holds (with appropriate modifications) for any left Artinian ring R with zero singular ideal such that \hat{R} is left Artinian. In particular the theorem holds whenever the injective envelope of ${}_R R$ is a finitely generated left R -module.

REFERENCES

1. CARTAN, H. AND EILENBERG, S. "Homological Algebra." Princeton University Press, Princeton, New Jersey, 1956.
2. GOLDIE, A. W. Torsion-free modules and rings, *J. A.*, 1 (1964), 268-287.
3. JOHNSON, R. E. Quotient rings of rings with zero singular ideal. *Pacific J. Math.* 11 (1961), 1385-1392.
4. LAMBEK, J. On Utumi's ring of quotients. *Canadian J. Math.* 15 (1963), 363-370.
5. MURASE, I. On the structure of generalized uniserial rings, I. *Sci. Papers, College General Education, Univ. Tokyo* 13 (1963), 1-22.